

**Ants (Hymenoptera: Formicidae) of the UC Davis Arboretum and the Häagen Dazs Honey Bee Haven,  
Yolo County, California**

Philip S. Ward  
Department of Entomology & Nematology  
University of California, Davis, CA 95616  
psward@ucdavis.edu

This list of ants from the UC Davis Arboretum, Yolo County, California (approximately 38°33'N 121°45'W) is based on observations made over a thirty-year period between 1985 and 2015. During this time there has been a steady decline in populations of native ants and an increasing dominance by the introduced Argentine ant, *Linepithema humile*. When this water-requiring species invades an area it is well known to eliminate most species of native ants (Ward, 1987; Holway, 1995; Suarez et al., 2001).

Subfamily, species	Nesting habits (A = arboreal) (G = ground)	Feeding habits (O = omnivore/scavenger) (P = predator) (S = seed-harvester)
<b>Ponerinae</b>		
<i>Hypoponera opacior</i>	G	P
<i>Hypoponera punctatissima</i>	G	P
<b>Myrmicinae</b>		
<i>Cardiocondyla mauritanica</i>	G	O
<i>Pheidole californica</i>	G	S/O
<i>Pogonomyrmex subdentatus</i>	G	S/O
<i>Solenopsis molesta</i>	G	O
<i>Solenopsis xyloni</i>	G	S/O
<i>Stenamma californicum</i>	G	P
<i>Stenamma punctatoventre</i>	G	P
<i>Stenamma</i> sp. CA02	G	P
<i>Tetramorium</i> sp. nr. <i>caespitum</i>	G	O
<b>Formicinae</b>		
<i>Brachymyrmex depilis</i>	G	O
<i>Camponotus essigi</i>	A	O
<i>Formica moki</i>	G	O
<i>Nylanderia vividula</i>	G	O
<i>Prenolepis imparis</i>	G	O
<b>Dolichoderinae</b>		
<i>Dorymyrmex bicolor</i>	G	O
<i>Dorymyrmex insanus</i>	G	O
<i>Liometopum occidentale</i>	A	O
<i>Linepithema humile</i>	A/G	O
<i>Tapinoma sessile</i>	G	O

Of the preceding 21 species, 16 are native and 5 are introduced taxa (*Hypoponera punctatissima*, *Cardiocondyla mauritanica*, *Tetramorium* sp. nr. *caespitum*, *Nylanderia vividula*, and *Linepithema humile*).

Ant species occurring in the vicinity of the Häagen Dazs Bee Garden (38°32'N 121°47'W) are highlighted.

The following two species also occur in or near the bee garden:

**Myrmicinae**

**Monomorium ergatogyna** G 0

**Formicinae**

**Formica aerata** G 0

**Descriptions of some ant species occurring in Davis**

Descriptions of some of the more conspicuous or interesting ant species of the UC Davis Arboretum are given below. Note that the body of a worker ant is composed of three major parts: (1) the head, (2) a middle section called the mesosoma, which corresponds roughly to the thorax, and (3) a posterior part called the gaster (roughly corresponding to the abdomen). At the constriction or "waist" between the mesosoma and the gaster is a node-like or scale-like structure called the petiole. In some species a second node, called the postpetiole, is interpolated between the petiole and the gaster. All descriptions below apply to the worker caste; the winged males and queens are often different in appearance.

**1. Liometopum occidentale (velvety tree ant)**

Small-medium, variable in size (3-6 mm long), orange-brown, with gray gaster. Mesosoma convex in lateral profile. Petiole scale-like, postpetiole absent.

*Liometopum occidentale* is an arboreal species that nests in cavities in the boles of large trees such as valley oak, walnut, and cottonwood. The ants are generalist predators and scavengers, and they frequently form large foraging columns, on tree trunks and on the ground. The workers have an unpleasant but distinctive (vinegary) odor. Various other native arthropods have co-evolved with this native ant: there are parasitic phorid flies that are host-specific and attack only the workers of *Liometopum*; there are mirid bugs that mimic the workers and often co-occur with them on tree trunks; and there are specialist staphylinid beetles that live in the ant nest. Thus, when *Liometopum occidentale* goes extinct as a result of invasion by the Argentine ant, we lose an entire community of associated arthropods.

**2. Pogonomyrmex subdentatus**

Medium-large (6-8 mm long), reddish-brown; postpetiole present. Sting present.

This is a large, red, seed-harvesting ant that nests in the ground. The nest entrances are usually decorated with small pebbles. Formerly more common on campus, this species survives along dry roadside verges, but does not tolerate the transformation to irrigated greenscapes.

**3. Formica moki (field ant)**

Medium-sized (6 mm long), with a dark head, orange-brown mesosoma and silvery-gray gaster. Mesosoma saddle-shaped in profile. Petiole scale-like, postpetiole absent.

This species was previously more widespread in the Arboretum. It nests under stones, in rotten wood, and directly in the soil. Workers are active foragers, and their movements are fast and skittish. They look superficially like workers of *Liometopum occidentale* but they are more uniform in size and less inclined to forage in large groups.

**4. Prenolepis imparis (winter honey ant)**

Small (3-4 mm long), brown, with shiny gaster. Mesosoma saddle-shaped in profile. Petiole scale-like; postpetiole lacking.

This is a common species, whose nests are located deep in the ground. The workers are quite cold-tolerant: they forage on the ground and on vegetation throughout the winter and spring months, and then cease above-ground activity during the summer drought. During this period of aestivation, the colony apparently subsists on accumulated food-stores, including honey stored in the swollen gasters of certain workers (called "repletes"). This is one of the very few native ants that can survive in areas invaded by the Argentine ant, probably because it is underground when the Argentine ants are most active.

#### 5. *Linepithema humile* (Argentine ant)

Small (2.5 mm long), grayish-brown, with a thin (scale-like) petiole; postpetiole absent.

This introduced species is especially common in disturbed and irrigated areas; forages in large files; and aggressively attacks and eliminates most native species of ants. Unlike most ant species, nests of Argentine ants generally exhibit no intraspecific aggression. In fact, there are not sharp colony boundaries—all the ants in a given geographical area behave as if members of one large supercolony.

### Literature Cited

Holway, D. 1995. Distribution of the Argentine ant (*Linepithema humile*) in northern California. *Conservation Biology* 9:1634-1637.

Suarez, A. V., Holway, D.A. & Case, T. J. 2001. Patterns of spread in biological invasions dominated by long-distance jump dispersal: insights from Argentine ants. *Proceedings of the National Academy of Sciences of the United States of America* 98:1095-1100.

Ward, P. S. 1987. Distribution of the introduced Argentine ant (*Iridomyrmex humilis*) in natural habitats of the lower Sacramento River Valley and its effects on the indigenous ant fauna. *Hilgardia* 55(2):1-16.

### Other Useful References on Ants

Bolton, B. 1994. *Identification guide to the ant genera of the world*. Cambridge, Mass.: Harvard University Press, 222 pp.

Bolton, B. 1995. *A new general catalogue of the ants of the world*. Cambridge, Mass.: Harvard University Press, 504 pp.

Fisher, B. L., Cover, S. P. 2007. *Ants of North America. A guide to genera*. Berkeley: University of California Press, xiv + 194 pp.

Hölldobler, B., Wilson, E. O. 1990. *The ants*. Cambridge, Mass.: Harvard University Press, 732 pp.

Hölldobler, B., Wilson, E. O. 2008. *The superorganism. The beauty, elegance, and strangeness of insect societies*. New York: W. W. Norton & Company.

Ward, P. S. 2005. A synoptic review of the ants of California (Hymenoptera: Formicidae). *Zootaxa* 936: 1-68. <http://www.mapress.com/zootaxa/2005f/zt00936.pdf>

Wheeler, G. C., Wheeler, J. 1963. *The ants of North Dakota*. Grand Forks, North Dakota: University of North Dakota Press, viii + 326 pp.

**Wheeler, G. C., Wheeler, J. 1973. Ants of Deep Canyon. Riverside, Calif.: University of California Press, xiii + 162 pp.**

**Wheeler, G. C., Wheeler, J. 1986. The ants of Nevada. Los Angeles: Natural History Museum of Los Angeles County, vii + 138 pp.**

#### **Web Sites**

**Ants of California: <http://www.antweb.org/california.jsp>**

**General information about ant diversity: <http://www.antweb.org>**

**Online catalog of the world's ants: <http://www.antcat.org/>**

**For additional (color) images of California ants see the "California ants" page of AntWeb: <http://www.antweb.org/california.jsp>**